# Intermittent Catheterisation Clinical Practice Principles

Diane K. Newman DNP, ANP-BC, FAAN, BCB-PMD; Angie Rantell PhD, ALNP; Piet Eelen RN, MSc, CNS-MS; Lisa Morgan MSN, RN, CNRN, AGCNS-BC; Morgan McDowell DNP, AGPCNP-BC, WHNP-BC; Rachel Kaminski BSN, RN, CWOCN, BCMAS



#### 1. Pre-treatment

Utilise current educational materials for IC and self-care

Ensure informed consent of the procedure, risks, benefits, and alternatives

Offer individualised instruction of catheter use



### Educate Collaborate Empower



## 2. Initial catheter selection

Provide a private, safe, dignified, and clean environment for learning about IC

Allocate adequate time for teaching

Offer information on various product choices and techniques<sup>1</sup>



## 3. The lifespan of intermittent catheter use

Patients should receive ongoing care and support

The opportunity to connect with external resources for support

Obtain supplies based on healthcare professional recommendations



Scan for additional resources and access to Convatec me+ Continence Care support or visit qr.convatec.com/cc-meplus-hcp

REFERENCES: 1.Burgess J, Gleba J, Lawrence K, Mueller S. Ostomy and Continent Diversion Patient Bill of Rights. Journal of Wound Ostomy Continence Nursing:2022;49(3):251-260. © 2024 Convatec. ™/® indicate trademarks of the Convatec group of companies. Convatec me+ is a trademark of Convatec Ltd. AP-70149-GBL-ENG-v1.